

Genetic diversity of mango leafhopper, *Amritodus atkinsoni* (Hemiptera: Cicadellidae) based on mtCOI gene sequences from India

Jaipal S. Choudhary, Chandra S. Prabhakar, Naiyar Naaz, Moanaro Lemtur, Bikash Das, Shivendra Kumar & Bhagwati P. Bhatt

To cite this article: Jaipal S. Choudhary, Chandra S. Prabhakar, Naiyar Naaz, Moanaro Lemtur, Bikash Das, Shivendra Kumar & Bhagwati P. Bhatt (2019) Genetic diversity of mango leafhopper, *Amritodus atkinsoni* (Hemiptera: Cicadellidae) based on mtCOI gene sequences from India, Mitochondrial DNA Part B, 4:1, 261-264, DOI: [10.1080/23802359.2018.1532332](https://doi.org/10.1080/23802359.2018.1532332)

To link to this article: <https://doi.org/10.1080/23802359.2018.1532332>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 23 Dec 2018.



Submit your article to this journal [↗](#)



Article views: 606



View related articles [↗](#)



View Crossmark data [↗](#)

Genetic diversity of mango leafhopper, *Amritodus atkinsoni* (Hemiptera: Cicadellidae) based on mtCOI gene sequences from India

Jaipal S. Choudhary^a, Chandra S. Prabhakar^{a,b} , Naiyar Naaz^a, Moanaro Lemtur^b, Bikash Das^a, Shivendra Kumar^a and Bhagwati P. Bhatt^c 

^aICAR Research Complex for Eastern Region Research Centre, Ranchi, India; ^bDepartment of Entomology, Veer Kunwar Singh College of Agriculture (Bihar Agricultural University, Sabour), Buxar, India; ^cICAR Research Complex for Eastern Region, ICAR Patna, Patna, India

ABSTRACT

The mango leafhopper, *Amritodus atkinsoni* is a serious and an endemic insect pest of mango in India. We analyzed mtCOI gene sequences from six Indian populations of *A. atkinsoni* for genetic diversity and population structure. The analysis of mtCOI sequence revealed 14 unique haplotypes and a low level of nucleotide diversity. mtCOI gene sequence analysis also revealed that *A. atkinsoni* specimens were clearly differentiated from other closely related species with a high level of accuracy.

ARTICLE HISTORY

Received 19 July 2018
Accepted 31 August 2018

KEYWORDS

Genetic structure;
haplotypes; phylogeography

Introduction

Mango leafhopper *Amritodus atkinsoni* (Lethierry) is one of the most economically important insect pests of mango (*Mangifera indica* L.) distributed mainly in South Asian and South East Asian countries (Waterhouse 1993). *Amritodus atkinsoni* is considered as the native of India and Pakistan (Pena et al. 1998) and causes 20–100% yield loss (Choudhary et al. 2012). In spite of the status of the pest, information on the population genetics of *A. atkinsoni* is not available from any region of the world.

Mitochondrial COI gene has been extensively used in population studies and identification of insects have unique features i.e. non recombinant, high copy numbers, simple maternal inheritance, high rate of evolution and robust evolutionary markers for determining intra- and inter-specific relationships of various invertebrate taxa including many insect species (Lunt et al. 1996; Armstrong and Ball 2005; Choudhary et al. 2018).

Therefore, in this study, we have employed the mitochondrial COI gene to infer the genetic diversity of *A. atkinsoni* collected from different regions of India. The dataset of mtCOI gene developed for the species has the potential for rapid and accurate identification.

Materials and methods

Adult *A. atkinsoni* were collected from six locations within India (Table 1 and Figure 1(A)). The collected adults were identified on the basis of morphological descriptions given by Viraktamath and Mohan (2004). Voucher specimens were

preserved with voucher ID: A1, A2, B1, B2, B3, B4 P3, P5, P8 and P6 in the Entomology Laboratory, ICAR RCER, Research Centre, Ranchi, India.

Total genomic DNA was extracted from individual specimens with methods described in Prabhakar et al. (2009). The mtCOI gene was amplified using UEA7 (F) and UEA10 (R) developed by Lunt et al. (1996) with PCR conditions mentioned in Prabhakar et al. (2012). These were then freeze-dried and custom sequenced using same upstream and downstream primers (Xcelris Labs Limited, India).

GenBank Accession Numbers: KC513467–KC513474 and KY084430–KY084442.

Nucleotide sequence analysis

The 85 mtCOI gene sequences of *A. atkinsoni* were aligned using ClustalW implemented in MEGA 6.0. (Tamura et al. 2013). Descriptive statistics of sequences were calculated with DnaSP (Librado and Rozas 2009). A median-joining network was constructed using NETWORK 4.6 (Bandelt et al. 1999). Phylogenetic reconstruction with 102 sequences representing the family Cicadellidae were analysed with the neighbor-joining method in MEGA 6.0. The “best close match” method in the programme TaxonDNA (Meier et al. 2006) was performed to test the frequency of successful identification.

Results and discussion

mtCOI gene sequences of 537 bp were obtained and aligned from 85 individuals of *A. atkinsoni* from six populations in India. The number of haplotypes per population (*H*) ranged

Table 1. Details of sampling locations and genetic diversity indices of *Amritodus atkinsoni*.

Location, State	ID code	Latitude (N)	Longitude (E)	Elevation m (amsl)	Collection year	n	H	Hd	K ± SD	π ± SD
Ranchi, Jharkhand	JH	23°35'	85°33'	651	2012	18	5	0.48366	0.69935 ± 0.55347	0.00131 ± 0.00115
SK Nagar, Gujarat	GU	24°12'	72°12'	209	2012	3	3	1.0	1.33333 ± 1.09834	0.00248 ± 0.002551
Sangareddy, Telangana	TL	17°62'	78°09'	524	2013	20	6	0.57368	0.76316 ± 0.58387	0.00142 ± 0.001214
Patna, Bihar	BH	25°61'	85°14'	58	2013	18	6	0.49020	0.77778 ± 0.59406	0.00145 ± 0.001237
Bhubaneswar, Odisha	OD	20°27'	85°84'	45	2013	17	3	0.22794	0.23529 ± 0.28567	0.00044 ± 0.000595
Jaipur, Rajasthan	RJ	26°92'	75°82'	431	2013	9	1	0.0	0.0 ± 0.0	0.0 ± 0.0
All						85	14	0.88291	0.641 ± 0.039	0.00164 ± 0.00019

n: Number of samples; H: Number of unique haplotypes; Hd: Haplotypes diversity; K: Average number of nucleotide differences; π: Nucleotide diversity; SD: Standard deviation.




Figure 1. (A) Sites of *Amritodus atkinsoni* collections for the present study. BH: Bihar, JH: Jharkhand, OD: Odisha, TL: Telangana, GU: Gujarat and RJ: Rajasthan. (B) Median joining network of mtCOI haplotypes of *Amritodus atkinsoni*. Each circle represents haplotype, and size of the circle is proportional to the haplotype frequency. Colours indicate the proportion of individuals of different populations present in a haplotype. (C) Neighbor-joining tree of 102 mtCOI gene sequences of different species representing family Cicadellidae. Bootstrap values based on 1000 replications are shown near nodes.

from 1 to 6, haplotype diversity (H_d) from 0 to 1.000, average number of nucleotide differences within populations (K) from 0 to 0.23529 ± 0.28567 and nucleotide diversity (π) ranged from 0 to 0.00248 ± 0.002551 (Table 1). A total of 14 haplotypes were identified. The most common haplotype was H2 and H5 comprised of 43 and 28 individuals of *A. atkinsoni*, respectively. Haplotype 2 (H2) was shared by five populations [JH (Jharkhand), GU (Gujarat), TL (Telangana), BH (Bihar) and OD (Odisha)] except RJ (Rajasthan) (Figure 1(B)). The analysis of mtCOI sequences demonstrated low values of nucleotide diversity and relatively high value of haplotype diversity with no geographic pattern. Most of the populations shared one or both of the most frequent haplotypes (H2 and H5). When values of haplotype ($H_d > 0.5$) and nucleotide ($\pi > 0.5\%$) diversity are high, the population appears to be stable with a relatively long evolutionary history (Rosetti and Remis 2012). None of the population presently examined meet these conditions, suggesting recent and unstable evolutionary history of the species. Low values of π and high values of H_d suggest the existence of small populations that have suffered recent population growth. Except R, this situation is observed for all the populations studied, where haplotype diversity is relatively high and nucleotide diversity relatively low. When both π and H_d values are low, the population has probably undergone a reduction in population size or a recent colonization event (Rosetti and Remis 2012). RJ is characterized by low values in both indices, reflecting recent population colonization.

The mtCOI gene analysis in the present study reveals *A. atkinsoni* to exhibit a high degree of diversity when compared with other hopper species. In *Scaphoideus titanus* Ball, COII gene sequences with 22 European and 8 Asian populations showed H_d and (π) value of 0.583 ± 0.072 & 0.203 ± 0.042 and 0.0046 ± 0.0027 & 0.0013 ± 0.0001 for nucleotide diversity, respectively, and showed moderate values for haplotype and nucleotide diversity (Papura et al. 2012). Genetic analysis of mtCOI gene sequences among 11 populations of rice brown planthopper, *Nilaparvata lugens* (Stål) and 5 populations of whitebacked planthopper, *Sogatella furcifera* (Horvath) from southeast Asia, showed low values of haplotype diversity 0.000–0.299 and 0.188–0.382 for *N. lugens* and *S. furcifera*, respectively (Mun et al. 1999), that were substantially lower than those observed in present study.

One hundred and two sequences from the family Cicadellidae were analysed for the accurate identification of *A. atkinsoni*. Intraspecific genetic divergence of 85 mtCOI gene sequences of *A. atkinsoni* ranged between 0 and 0.9% with an average of 0.50%. The best close match methods identified the 95% intraspecific genetic divergence threshold value 0.44% of *A. atkinsoni* identified all *A. atkinsoni* specimens correctly. Our results showed large gaps between intra and interspecific genetic distances for cicadellid species are consistent with previous reports on mtCOI gene based identification of insect species (Choudhary et al. 2018). DNA species barcode gaps can be detected by recording the overlap between the highest intraspecific and the lowest interspecific genetic distances (Meier et al. 2008). Phylogenetic analysis using the neighbor-joining method (Figure 1(C)) gave a

strong bootstrap support (99%) for the monophyly of *A. atkinsoni*. The mtCOI gene sequences developed in the present study and its availability in GenBank could also be useful in identification of the species in the future (Virgilio et al. 2010; Jinbo et al. 2011). Incorrect species identification using the mtCOI is largely down to insufficient geographic sampling of the taxa (Jinbo et al. 2011). In this context and in relation to *A. atkinsoni*, our present results significantly increase the number of mtCOI gene sequences available, which will encourage the molecular identification of this and related important mango leafhopper species.

Acknowledgements

Authors are grateful to Mr. Deepak Tanti and Mr. Pradeep Lakra for collection of mango leafhoppers.

Disclosure statement

The authors declare no conflicts of interest.

Funding

This work was supported by the Indian Council of Agricultural Research (ICAR), New Delhi, for providing financial assistance for this study under the National Innovations in Climate Resilient Agriculture (ICAR-RCER/RC R/E.F./2011/29).

ORCID

Chandra S. Prabhakar  <http://orcid.org/0000-0001-7182-2667>
Bhagwati P. Bhatt  <http://orcid.org/0000-0002-2575-6619>

References

- Armstrong KF, Ball SL. 2005. DNA barcodes for biosecurity: invasive species identification. *Philos Trans R Soc Lond B*. 360:1813–1823.
- Bandelt HJ, Forster P, Rohl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol*. 16:37–48.
- Choudhary JS, Naaz N, Lemtur M, Das B, Singh AK, Bhatt BP, Prabhakar CS. 2018. Genetic analysis of *Bactrocera zonata* (Diptera: Tephritidae) populations from India based on *cox1* and *nad1* gene sequences. *Mitochondr DNA A*. 29:727–736. DOI: [10.1080/24701394.2017.1350952](https://doi.org/10.1080/24701394.2017.1350952)
- Choudhary JS, Prabhakar CS, Maurya S, Kumar R, Das B, Kumar S. 2012. New report of *Hirsutella* sp. infecting mango hopper *Idioscopus clypealis* from Chotanagpur Plateau, India. *Phytoparasitica*. 40:243–245.
- Jinbo U, Kato T, Ito M. 2011. Current progress in DNA barcoding and future implications for entomology. *Entomol Sci*. 14:107–124.
- Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*. 25:1451–1452.
- Lunt DH, Zhang DX, Szymura JM, Hewitt GM. 1996. The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies. *Insect Mol Biol*. 5:153–165.
- Meier R, Shiyang K, Vaidya G, Ng PKL. 2006. DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Syst Biol*. 55:715–728.
- Meier R, Zhang G, Ali F. 2008. The use of mean instead of smallest interspecific distances exaggerates the size of the “barcode gap” and leads to misidentification. *Syst Biol*. 57:809–813.
- Mun J, Song Y, Heong KL, Roderick GK. 1999. Genetic variation among Asian populations of rice planthoppers, *Nilaparvata lugens* and *Sogatella furcifera* (Hemiptera: Delphacidae): mitochondrial DNA sequences. *Bull Entomol Res*. 89:245–253.

- Papura D, Burban C, van Helden M, Giresse X, Nusillard B, Guillemaud T, Kerdelhué C. 2012. Microsatellite and mitochondrial data provide evidence for a single major introduction for the Nearctic Leafhopper *Scaphoideus titanus* in Europe. PLoS One. 7:e36882.
- Pena JE, Mohyuddin AI, Wysoki M. 1998. A review of the pest management situation in mango agroecosystems. Phytoparasitica. 26:129–148.
- Prabhakar CS, Mehta PK, Sood P, Singh SK, Sharma P, Sharma PN. 2012. Population genetic structure of the melon fly, *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae) based on mitochondrial *cytochrome oxidase* (COI) gene sequences. Genetica. 140:83–91.
- Prabhakar CS, Sood P, Kapoor V, Kanwar SS, Mehta PK, Sharma PN. 2009. Molecular and biochemical characterization of three bacterial symbionts of fruit fly, *Bactrocera tau* (Tephritidae: Diptera). J Gen App Microbiol. 55:213–220.
- Rosetti N, Remis MI. 2012. Spatial Genetic Structure and Mitochondrial DNA Phylogeography of Argentinean Populations of the Grasshopper *Dichroplus elongatus*. PLoS one. 7:e40807.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. Mol Biol Evol. 30: 2725–2729.
- Viraktamath CA, Mohan GS. 2004. A revision of the *Idiocerine* leafhopper genus *Amritodus* (Hemiptera: Cicadellidae) breeding on mango. Entomon. 578:1–117.
- Virgilio M, Backeljau T, Nevado B, Meyer MD. 2010. Comparative performances of DNA barcoding across insect orders. BMC Bioinformatics. 11: 206.
- Waterhouse DF. 1993. The major arthropods pests and weeds of agriculture in South-East Asia; Distribution, Importance and Origin. Monograph. No. 21.